

## Actero™ Buffered Listeria Enrichment Broth Product Information

Catalogue No.	Description
<b>FCM-057</b>	Actero™ Buffered Listeria Enrichment Broth (500 G)
<b>FCM-056</b>	Actero™ Buffered Listeria Enrichment Broth (2 KG)
<b>FCM-055</b>	Actero™ Buffered Listeria Enrichment Broth (10 KG)

**INTENDED USE:**

Buffered Listeria Enrichment Broth is used for selective enrichment of *Listeria* spp. A modification of Listeria Enrichment Broth developed by Lovett, this media was designed to improve the enrichment of *Listeria* spp. with the addition of Disodium Phosphate thereby increasing the buffering capacity. *Listeria* spp., found in soil, sewage, and river water in addition to unprocessed foods, are Gram-positive, short and motile rods. Cycloheximide in this formulation inhibits the growth of saprophytic fungi while Nalidixic Acid is used to inhibit the growth of Gram-negative organisms.

**Formula\* per Liter:**

Casein Digest of Peptone.....	17.0g
Soy Peptone.....	3.0g
Dextrose.....	2.5g
Yeast Extract.....	6.0g
Nalidixic Acid.....	0.04g
Cycloheximide.....	0.05g
Monopotassium Phosphate.....	1.35g
Sodium Chloride.....	5.0g
Acriflavine.....	0.015g
Dipotassium Phosphate.....	2.5g
Disodium Phosphate.....	9.6g

**Final pH: 7.3 ± 0.2 at 25°C**

\* Grams per liter may be adjusted or formula supplemented to obtain the desired performance.

**PREPARATION:**

Mix 47 grams of the medium in one liter of purified water until evenly dispersed. Heat with repeated stirring and boil for one minute to dissolve completely. Distribute and autoclave at 121°C for 15 minutes.

**TERMS AND CONDITIONS:**

Salus Scientific Inc. makes no representations and warranties concerning its products other than those stated herein. All Product(s) delivered hereunder by Salus Scientific Inc., its affiliates or any other person on its behalf shall, at the time of delivery, be manufactured to meet Salus Scientific Inc.'s specifications and all applicable laws.

All other terms, conditions and warranties, including any warranty of merchantability, quality, fitness or suitability for a particular or intended purpose, implied by common law or statute, (implied warranties) are expressly excluded.

**FOR FURTHER INFORMATION, PLEASE CONTACT:**

Salus Scientific Inc.  
Suite 450, 1414 – 8 St. S.W.  
Calgary, Alberta, Canada  
T2R 1J6  
Tel: 1-877-298-0208

**QUALITY CONTROL SPECIFICATIONS:**

1. The powder is homogeneous, free-flowing, and light beige to beige.
2. Visually the prepared medium is gold to yellow, and slightly hazy to clear.
3. Expected cultural response after 18-48 hours at 30 ± 2°C.

**Result:**

Organism	Result
<i>Escherichia coli</i> ATCC® 25922	Inhibited
<i>Listeria monocytogenes</i> ATCC® 7644	Growth
<i>Staphylococcus aureus</i> ATCC® 25923	Suppressed 18-24 hours
<i>Listeria monocytogenes</i> ATCC® 15313	Growth

**Storage Instructions:**

Store the sealed bottle containing the dehydrated medium at 2 to 30°C. Once opened and recapped, place the container in a low humidity environment at the same storage temperature. Protect it from moisture and light. The dehydrated medium should be discarded if it is not free flowing, or if the color has changed from the original light beige to beige color.